

Scientific Contributions (cont'd.)

Effect of supplementation with magnesium and tryptophan on the welfare and meat quality of pigs

Núria Panella, Maria Font i Furnols, Ma. Àngels Oliver, Pedro Rodríguez, Antonio Velarde, Joan Tibau¹ and Marta Gil

IRTA. Centre de Tecnologia de la Carn. Granja Camps i Armet. 17121 Monells. Girona. Spain.

¹*IRTA. Centre de Control Porcí. Veïnat de Sies. 17121 Monells. Girona. Spain.*

Background

Handling of animals during transport, lairage and slaughter is one of the most important causes of stress in pigs and that, together with genotype, most influence the final carcass and meat quality. A supplementation of magnesium (Mg) in the diet of pigs may reduce the response of the animals to stressful stimuli by reducing the levels of cortisol and catecholamines in plasma. An increase of tryptophan (Trp) concentration in the diet enhances the synthesis of brain serotonin and may have sedative effects. Some authors report an improvement of pork meat quality as a result of the supplementation of Mg or Trp. However, other studies contradict these findings or obtain inconsistent results. The effects of the supplement of Mg on meat quality seem to be related to the stress-susceptibility of the pigs, that is to their halothane genotype. On the other hand, in an Australian study it was found that under a negative antemortem handling treatment, a dietary Mg supplementation significantly improves pork quality and reduces the incidence of PSE carcasses. Accordingly, a Dutch study reported that a short-term supplementation of the diet with magnesium acetate or with a combination of this salt with Trp and

vitamins does not improve pork quality when pigs are not stressed beyond levels associated with routine slaughter procedures.

Objectives

The aim of this study was to evaluate the effect of the supplementation of the diet with Mg and Trp on the response to stressful factors and on meat quality of pigs. Animals homozygous with respect to the halothane gene (positive, nn, and negative, NN) were used to achieve this objective.

Materials and methods

Animals, diets and pre-slaughter treatment: Seventy-one entire male pigs (36 NN and 35 nn) from Pietrain, Landrace and Large White lines, with an average live weight of 108.51 ± 8.32 kg, were used in this study. All animals were housed individually in pens with a space allowance of 4 m², permitting visual and olfactory contact between each other. They were fed the same diet until 5 days before slaughtering, when three diet groups were established for each genotype: diet 1 group, same diet supplemented with 1.2 g elemental magnesium (Mg) and 8 g L-tryptophan (Trp) per kg; diet 2 group, same diet supplemented with 8 g Trp per kg, and control group, with no supplement. Two batches of 35 and 36 animals were slaughtered on 2 different days. The animals of each batch were mixed in 6 different groups according to diet and genotype and transported from the farm to the experimental abbatoir for one hour on rougher secondary roads. Lairage time for the first pigs slaughtered was about 30 min and for the last ones was of 7 h. They were all stunned with 90% CO₂ and slaughtered. In each batch, the animals from the 6 groups were slaughtered alternatively. Hot carcass weight was measured at 45 min. *postmortem* (pm), and used to calculate the killing-out (%).

Skin lesions: After sticking, and when the carcasses were hoisted on the bleeding rail, fresh scratches on the skin

Scientific Contributions (cont'd.)

were inspected and subjectively scored according to their location (head/neck, flanks/back and hindquarter) and occurrence (0 = No occurrence, 1 = 1 occurrence, 2 = 2 to 5 occurrence, and 3 = more than 5).

Meat quality measurements: The left side of the carcass was used to perform meat quality measurements on the *Longissimus thoracis* (LT) and *Semimembranosus* (SM) muscles. Muscle pH at 45 min (pH45) and at 24 h (pHu) pm were measured using a portable pH meter equipped with a Xerolyt electrode. Electrical conductivity at 24 h pm (ECu) was measured at the last rib level using a Pork Quality Meater. The muscle water holding capacity (WHC) was determined in the LT muscle using the bag method.

Colour measurements: Colour measurements were carried out with a Minolta Chroma Meter CR-200 using the white tile provided by the manufacturer as the internal standard and set to illuminant C. Triplicate measurements were taken on LT muscle. The measurements were expressed as CIE $L^* a^* b^*$.

Results and discussion

Five animals died before being slaughtered (during transport or while unloading the lorry), all of them from halothane positive pigs (nn): three corresponded to the control and the other two to the diet 1 group (Mg + Trp). The sedative effect of the diet with Trp (diet 2) seemed to facilitate the coping of the animals to stressful factors, such as new environments. The presence and severity of skin lesions indicative of aggressive behaviour are presented in Table 1. Skin lesions in the head/neck, flanks/back and hindquarters were significantly higher in the group of animals fed diet 1 compared with the animals of the control group. No differences were found in the occurrence of skin lesions between diet 1 and diet 2, except in the flanks. The increase in skin lesions of the animals of diet 1 and 2 would suggest an increase of activity due to fighting and a depletion of the glycogen resource before slaughter. This was

reflected in the ultimate pH and other meat quality variables. Table 2 shows the effect of the diet on several meat quality variables: pHu SM tended to be higher ($P=0.080$) in the animals fed diet 1 and diet 2 than in the control. The incidence of DFD carcasses (dark, firm and dry), established as pHu SM ≥ 6.0 , was 12.5% in the diet 1 group and 16.7% in the diet 2 group, and 0% in the control. The ECu LT tended to be slightly lower ($P = 0.09$) when animals were fed diet 2. The most important differences among groups were found for the L^* (lightness) and the drip loss variables. L^* values were lower ($P < 0.05$), indicating darker muscle colour, in the LT of pigs fed the supplemented diets than in the LT of controls, whereas drip losses were lower ($P < 0.05$) in the LT of animals fed with diets 1 and 2 than the control. Thus, the supplementation with Mg+Trp or with Trp improved the muscle colour and the WHC. From the present preliminary results and in the conditions of this experiment, it would seem that the darker colour and the decrease in drip losses of the muscle were a consequence of the inclusion of a supplement of (Mg+Trp) and Trp in the diet.

Conclusions

The mixture of Mg and Trp increased the frequency of skin lesions due to fighting during transport and lairage. The aggressive behaviors due to mixing contribute to an increase in pHu in the meat from the animals fed diet 1 and diet 2 and, as a consequence, some meat quality characteristics (colour and water holding capacity) improved. However, to ascertain if these supplements are adequate to be used in animal production in order to improve meat quality, other factors such as food intake and daily growth should be considered.

Acknowledgements

This work was financed by CICYT AGL2002-00777. The authors thank the Associacio Catalana de Criadors de Bestiar Porci Selecte - that includes 8 pig breeders - for

Scientific Contributions (cont'd.)

the availability of the pigs evaluated in this trial.

References

Adeola O., Ball R.O. (1992) Hypothalamic neurotransmitter concentrations and meat quality in stressed pigs offered excess dietary tryptophan and tyrosine. *J. Anim. Sci.* 70, 1888-1894.

Apple J.K., Maxwell C.V., deRodas B., Watson H.B., Johnson Z.B. (2000) Effect of magnesium mica on performance and carcass quality of growing-finishing swine. *J. Anim. Sci.* 78, 2135-2143.

Apple J.K., Maxwell C.V., Stivarius M.R., Rakes L.K., Johnson Z.B. (2002) Effects of dietary magnesium and halothane genotype on performance and carcass traits of growing-finishing swine. *Livest. Prod. Sci.* 76, 103-113.

Caine W.R., Schaefer A.L., Aalhus J.L., Dugan M.E.R. (2000) Behaviour, growth performance and pork quality of pigs differing in porcine stress syndrome genotype receiving dietary magnesium aspartate hydrochloride. *Can. J. Anim. Sci.* 80, 175-182.

CIE. (1976). Commission Internationale de l'Éclairage. Colorimetry. (Vol. Publication n° 15). Vienna, Austria: Bureau Central de la CIE.

D'Souza D.N., Warner R.D., Dunshea F.R., Leury B.J. (1999) Comparison of different dietary magnesium supplements on pork quality. *Meat Sci.* 51, 221-225.

D'Souza D.N., Warner R.D., Leury B.J., Dunshea F.R. (1998) The effect of dietary magnesium aspartate supplementation on pork quality. *J. Anim. Sci.* 76, 104-109.

D'Souza D.N., Warner R.D., Leury B.J., Dunshea F.R. (2000) The influence of dietary magnesium supplement type, and supplementation dose and duration, on pork quality and the incidence of PSE pork. *Aust. J. Agric. Res.* 51, 185-189.

Frederick, B.R., van Heugten, E., See, M.T. (2004) Timing of magnesium supplementation administered through drinking water to improve fresh and stored pork quality. *J. Anim. Sci.* 82, 1454-1460.

Geesink G.H., van Buren R.G.C., Savenije B., Verstegen M.W.A., Ducro B.J., van der Palen

J.G.P., Hemke, G. (2004) Short-term strategies and pork quality. *Meat Sci.* 67, 1-6.

Hamilton D.N., Ellis M., Hemann M.D., McKeith F.K., Miller K.D., Purser K.W. (2002) The impact of longissimus glycolytic potential and short-term feeding of magnesium sulfate heptahydrate prior to slaughter on carcass characteristics and pork quality. *J. Anim. Sci.* 80, 1586-1592.

Hamilton D.N., Ellis M., McKeith F.K., Eggert J.M. (2003) Effect of level, source, and time of feeding prior to slaughter of supplementary dietary magnesium on pork quality. *Meat Sci.* 65, 853-857.

Henry Y., Sève B., Mounier A., Ganier P. (1996) Growth performance and brain neurotransmitters in pigs affected by tryptophan, protein and sex. *J. Anim. Sci.* 74, 2700-2710.

Honikel, O.K. (1997). Reference methods supported by OECD and their use in Mediterranean meat products. *Food Chem.* 59, 573-582.

Kietzmann M & Jablonski H (1985) Blocking of stress in swine with magnesium aspartate hydrochloride. *Prakt. Tierz.* 661, 331.

Leathwood, P.D. (1987) Tryptophan availability and serotonin synthesis. *Proc. Nutr. Soc.* 46, 143

O'Quinn P.R., Nelssen J.L., Unruh J.A., Goodband R.D., Woodworth J.C., Tokach M.D. (2000) Effects of feeding modified tall oil and supplemental potassium and magnesium on growth performance, carcass characteristics, and meat quality of growing-finishing pigs. *Can. J. Anim. Sci.* 80, 443-449.

Schaefer A.L., Murray A.C., Tong A.K.W., Jones S.D.M., Sather A.P. (1993) The effect of ante mortem electrolyte therapy on animal physiology and meat quality in pigs segregating at the halothane gene. *Can. J. Anim. Sci.* 73, 231-240.

Swigert K.S., Mc Keith F.K., Carr T.C., Brewer M.S., Culbertson M. (2004) Effects of dietary vitamin D3, vitamin E, and magnesium supplementation on pork quality. *Meat Sci.* 67, 81-86.

Scientific Contributions (cont'd.)

*This paper has been previously published in the
Proceedings of the 50th International Congress
of Meat Science and Technology, 8-13th
August, 2004, Helsinki, Finland*

Table 1. Least squares means and standard errors of skin lesions mean values in pigs from three groups of diets

	CONTROL (n = 19)		DIET 1* (n = 22)		DIET 2* (n = 23)	
	LSM	S.E.	LSM	S.E.	LSM	S.E.
Head/neck	0.93 ^b	0.195	1.58 ^a	0.180	1.17 ^{ab}	0.176
Flanks/back	0.82 ^b	0.205	1.80 ^a	0.190	1.17 ^{ab}	0.186
Hindquarters	0.34 ^b	0.161	1.06 ^a	0.149	0.83 ^a	0.146

^{a,b} Different superscripts within a row indicated significant differences ($P < 0.05$).

*DIET 1: Magnesium and Tryptophan; DIET 2: Tryptophan.

(0 = No occurrence; 1 = 1 occurrence; 2 = 2 to 5 occurrence; and 3 = more than 5).

Table 2. Least squares means and standard errors of meat quality variables in the LT and SM muscles of pigs from the two genotypes fed the three groups of diets

	CONTROL (n = 20)		DIET 1* (n = 22)		DIET 2* (n = 24)	
	LSM	S.E.	LSM	S.E.	LSM	S.E.
Killing out (%)	77.67	0.391	78.22	0.371	77.47	0.355
Carcass weight at 24 h (kg)	43.18	0.580	41.95	0.536	42.04	0.5151
pH45 LT	6.01	0.059	6.04	0.055	6.10	0.053
pH45 SM	6.12	0.071	6.18	0.066	6.10	0.063
pHu LT	5.47	0.043	5.53	0.041	5.58	0.039
pHu SM	5.59	0.065	5.79	0.062	5.67	0.059
ECu LT	6.82	0.409	6.84	0.388	5.79	0.371
ECu SM	8.16	0.513	7.49	0.487	7.18	0.466
L* (lightness)	54.06 ^a	0.999	50.82 ^b	0.948	51.04 ^b	0.907
A* (redness)	0.79	0.231	1.22	0.214	1.22	0.206
B* (yellowness)	5.10	0.394	4.45	0.374	4.41	0.357
WHC (%)	8.72 ^a	0.460	7.34 ^b	0.436	7.35 ^b	0.417

^{a,b} Different superscripts within a row indicated significant differences ($P < 0.05$).

*DIET 1: Magnesium and Tryptophan; DIET 2: Tryptophan.